

Is it thrombotic microangiopathy (TMA)?

Consider TMA in lupus patients with thrombocytopenia, microangiopathic hemolytic anemia, and organ involvement¹⁻⁵



TMA can be associated with numerous conditions and triggers¹

TMA in patients with lupus—it could be atypical-HUS

If TMA is suspected, time is of the essence

Definition of TMA^{1,2}




Plus 1 or more of the following signs and symptoms:

Organ manifestations

Common signs and symptoms




Other



Evaluate ADAMTS13 activity and Shiga toxin/EHEC test^a

While ADAMTS13 results are awaited, a platelet count >30 x 10⁹/L and/or sCr >1.7 to 2.3 mg/dL almost eliminates a diagnosis of severe ADAMTS13 deficiency (TTP)



^aShiga toxin/EHEC test is warranted with history/presence of GI symptoms. ^bRange found in published literature is <5%-10%. ^cWhere TTP and STEC-HUS have been ruled out, the TMA can be attributed to complement dysregulation. Atypical-HUS is a complement-mediated TMA (CM-TMA) that has been more broadly or narrowly defined in different contexts, based on such factors as the presence and nature of a trigger and/or identification of an underlying genetic mutation. ^dMany other conditions may trigger atypical-HUS.

Atypical-HUS:

- is a serious, life-threatening medical disorder that can manifest with or be triggered^d by SLE^{1,3,6-9}
- is associated with complement dysregulation that leads to microvascular thrombosis, hemolysis, and ischemic organ injury^{1,2,10,11}
- may be a complication of lupus^{2,7,9}

Differential diagnosis of TMA ^{1,5,10-12}
SIGNS OF TMA
- Platelet count, blood smear showing schistocytes, signs of organ involvement, elevated LDH
TRIGGERS OF TMA
- Autoimmune disorders (eg, SLE, catastrophic antiphospholipid syndrome) Associated tests: antiphospholipid, LAC
- Cobalamin C deficiency Associated tests: B12, homocysteine, methylmalonic acid/methionine testing, <i>MMACHC</i> mutation screening
- DIC Associated tests: PT/aPTT, D-dimer
- Infection Associated tests: Shiga toxin panel, COVID-19, other infection panels
- TTP Associated test: ADAMTS13 activity level
- Other triggers (eg, pregnancy, genetic mutation)
- Atypical-HUS There is no single definitive test. Diagnosis is exclusionary

ADAMTS13=a disintegrin and metalloproteinase with a thrombospondin type 1 motif member 13; CV=cardiovascular; DIC=disseminated intravascular coagulation; EHEC=enterohemorrhagic *Escherichia coli*; GI=gastrointestinal; HUS=hemolytic uremic syndrome; LAC=lupus anticoagulant; LDH=lactate dehydrogenase; *MMACHC*=methylmalonic aciduria and homocystinuria type C; PT/aPTT=prothrombin time/activated partial thromboplastin time; sCr=serum creatinine; SLE=systemic lupus erythematosus; STEC=Shiga toxin-producing *Escherichia coli*; TMA=thrombotic microangiopathy; TTP=thrombotic thrombocytopenic purpura.

In a patient with lupus and thrombotic microangiopathy, consider atypical-HUS



In patients with lupus and persistent TMA, a timely diagnosis of atypical-HUS can help with necessary and specialized atypical-HUS disease management.^{1,13}

Learn more about TMA and atypical-HUS at
aHUSSource.com/physician

References

1. Azoulay E, et al. *Chest*. 2017;152(2):424-434.
2. Laurence J, et al. *Clin Adv Hematol Oncol*. 2016;14(11)(suppl 11):2-15.
3. Chen MH, et al. *Rheumatology (Oxford)*. 2011;50(4):768-775.
4. Pivovarova AI, et al. *Diseases*. 2020;9(1):3.
5. Shen YM. *Thromb J*. 2016;14(suppl 1):19.
6. Samson M, et al. *Intern Med J*. 2012;42(1):95-98.
7. Kello N, et al. *Semin Arthritis Rheum*. 2019;49(1):74-83.
8. Asif A, et al. *J Nephrol*. 2017;30(3):347-362.
9. Song D, et al. *Arthritis Res Ther*. 2013;15(1):R12.
10. Noris M, et al. *Nat Rev Nephrol*. 2012;8(11):622-633.
11. Brocklebank V, et al. *Clin J Am Soc Nephrol*. 2018;13(2):300-317.
12. Petri M. *Hematology Am Soc Hematol Educ Program*. 2019;2019(1):415-420.
13. Ono M, et al. *Intern Med*. 2018;57(11):1617-1623.